

3.7. The benthic communities of the Cap de Creus canyon

Teresa Madurell¹, Covadonga Orejas^{1,2}, Susana Requena¹, Andrea Gori^{1,3}, Ariadna Purroy¹, Claudio Lo Iacono⁴, Ana Sabatés¹, Carlos Dominguez-Carrió¹, Josep Maria Gili¹

¹ Institut de Ciències del Mar, ICM-CSIC, 08003 Barcelona, Spain

² Present address: Instituto Español de Oceanografía, IEO, 39004 Santander, Spain

³ Present address: Centre Scientifique de Monaco, 98000 Monaco

⁴ Unitat de Tecnologia Marina, UTM-CSIC, 08003 Barcelona, Spain

CAP DE CREUS CANYON: LOCATION AND ABIOTIC CHARACTERISTICS

The Cap de Creus canyon is located at the westernmost part of the Gulf of Lion continental margin, considered to be the last of a dense network of submarine canyons that characterize the area (Fig. 1). The head of the canyon is located about 5 km north-east from the Cap de Creus promontory and its end side meets with the end of the Sète Canyon at around 2150 m depth. The Cap de Creus canyon incises the shelf edge at depths of 110-130 m for more than 50 m and progressively widens to attain a maximum amplitude of 6 km, where it reaches depths ranging from 650 m to 2200 m. Both flanks of the canyon show characteristic morphological features due to differential depositional settings and hydrodynamic regimes. While the northern side displays a smooth morphology with rounded gullies and scars, showing a sedimentary depositional set of conditions, the southern flank has an erosive sedimentary nature, characterized by extensive rocky outcrops, vertical walls and terraces (Lastras *et al.*, 2007; DeGeest *et al.*, 2008; Puig *et al.*, 2008). Extensive linear erosive furrows occur at depths between 150 and 1400 m which appear to be associated with the dense-shelf water cascading phenomenon and represent the preferential routes for sediment transport along the southern flank of the canyon (Canals *et al.*, 2006; Lastras *et al.*, 2007; Puig *et al.*, 2008; Ulses *et al.*, 2008).

The proximity of the canyon to the coast, and therefore the reduced extension of the adjacent continental shelf, is another remarkable feature of this canyon. The continental shelf extends to about 130 m depth, displays a rough morphology and is very narrow near the Cap de Creus promontory and the canyon (2.7 km). The proximity to the coast added to the global configuration regarding the Gulf of Lion renders this canyon very particular. The Cap de Creus canyon appears to be a major export path of particulate matter (coming from river input and seasonal storms) from the shelf to the deep areas. The preferential direction of the coastal currents and the

narrowing of the shelf towards the west result in the majority of off-shelf sediment transport on this margin running through the Cap de Creus canyon. Sediment fluxes observed in this canyon are much higher than in eastern and central Mediterranean submarine canyons (Palanques *et al.*, 2006). Dense-shelf water cascading, which occurs more frequently and with higher intensity in this canyon, also favors the supply of nutrients and oxygen to the deeper parts of the canyon (Durrieu de Madron *et al.*, 2005; Canals *et al.*, 2006). This process has important biological and ecological implications that contribute to make it a unique system. The large amounts of organic material transported along the Cap de Creus canyon probably play an essential role in the maintenance of high local biodiversity rates and its associated deep-sea ecosystems (Canals *et al.*, 2009; Orejas *et al.*, 2009).




Fig. 1:

A) Cap de Creus canyon in the North Western Mediterranean context and B) its digital elevation model (DEM).

BENTHIC COMMUNITIES IN THE CANYON

The benthic communities found at the head of the canyon and its neighbouring area of the continental shelf have been studied for the last ten years and are still the subject of thorough research studies. The different types of megabenthic communities that dwell in this region are determined by the geological and environmental configuration of the continental shelf and canyon. Two main types of benthic communities have been identified in the southern flank of the canyon between 150 and 400 m depth: circalittoral communities also found on the continental shelf, and bathyal communities of cold-water corals (Gili *et al.*, 2011; Lo Iacono *et al.*, 2012). The description of the megabenthic communities is made after Pérès and Picard (1964), Desbruyères *et al.* (1972–1973) and Pérès (1985).

Circalittoral communities

The circalittoral communities identified are located on the shelf edge and break where sediments are very clean of silt and gravel predominates. There are two communities defined here:

1. The offshore detritic community of the shelf edge and break ("*biocoenose des fonds détritiques du large*") dominates in the outer shelf region (estimated surface area of 31.95%) (Lo Iacono *et al.*, 2012). It is characterized by sessile macrofauna, mainly the sea pens *Pteroeides spinosum*, *Pennatula rubra* and *Cavernularia pusilla*, together with the soft coral *Alcyonium palmatum* and sponges. Two different *facies* can be distinguished in the upper canyon rim and canyon head, those with shared dominance between sea pens and ceriantharians, and those dominated by sea pens and hydrozoans (Fig. 2 A, B).




Fig. 2:

Schematic representation of the offshore detritic communities dominated by (A) sea pens and ceriantharians and (B) sea pens and hydrozoans.

2. The offshore rocky-bottom community at the end of the continental shelf (outer shelf) (Fig. 3), known as the "*biocoenose de la roche du large*", has an estimated surface area of 1.81% (Lo Iacono *et al.*, 2012). This community is characterized by the presence of relatively small superficial rocky outcrops that are colonized by a rich fauna of sessile organisms, dominated by sponges like *Phakellia ventilabrum* and *Poecillastra compressa*; colonies of the cnidarian *Eunicella sp.*, and a variety of hydroids, bryozoans and ophiurids (genus *Ophiura*). Several species characteristic of both deep areas of the continental shelf as well as individuals from shallower coralligenous communities (e.g. *Parazoanthus axinellae* or *Dysidea tufa*) are also found in well-preserved rocky areas (Gili *et al.*, 2011). This community has been affected by trawl fishing and its presence in the Cap de Creus Canyon is remarkable.

Cold-water coral community

The cold-water coral community found on the southern wall of the Cap de Creus canyon is known as the "*biocoenose des coraux blancs*". This community develops on rocky substrates, dominated by boulders and vertical walls (Fig. 4). The most abundant cold-water coral species present in the Cap de Creus canyon are *Madrepora oculata*, *Lophelia pertusa*, *Dendrophyllia cornigera* and *Desmophyllum dianthus* (known in the Mediterranean as *D. cristagalli*) (Fig. 5), the dominant species being *M. oculata*. Quantitative analysis of occurrence and density of *M. oculata* shows a preferential presence of this species in rocky substrates and a spatial distribution pattern in patches of varying size reaching maximum densities of around 11 colonies/m² (Orejas *et al.*, 2009). *D. cornigera* has always been found either alone or in small groups, and *Lophelia pertusa* as isolated colonies.

Another feature of the cold-water coral community in the Cap de Creus submarine canyon is the high diversity of associated species, mainly of benthic suspension feeders such as brachiopods (especially *Gryphus vitreus* and *Mergella truncata*), polychaetes (*Sabella pavonina*), sponges, bryozoans and other cnidarians including, for instance, octocorals like the red coral *Corallium rubrum*.

The cold-water coral occurrence in the rocky areas of the southern wall of the Cap de Creus canyon is probably related to the food supply through the energetic current flows that periodically carry nutritive suspended particles from shelf environments (Canals *et al.*, 2006) and through the resuspension of shelf organic matter by inertial internal waves (Ogston *et al.*, 2008; Orejas *et al.*, 2009; Tsounis *et al.*, 2010; Purser *et al.*, 2010). The presence of corals in available hard substrates is also related to the reduced sediment accumulation rates caused by cascading phenomena and in general by the strong currents found in the area (DeGeest *et al.*, 2008; Puig *et al.*, 2008). The canyon's northern flank is mainly characterized by soft sediments with high accumulation rates (DeGeest *et al.*, 2008). The lack of rocky substrates probably explains the scarce presence of cold-water corals on the northern wall. However, explorations carried out in this flank of the canyon have been punctual and consequently the communities have not been thoroughly studied yet. Further surveys in the Cap de Creus canyon will provide a comprehensive map of the habitats present in the whole study area.




Fig. 3:
Detail of the rocky-bottom community found at the deeper parts of the continental shelf and canyon head, dominated by sponges.



Fig. 4:
Colonies of *Madrepora oculata* found at depths of 250 m on the southern flank of the Cap de Creus canyon.




Fig. 5:
Detail of the four dominant cold-water corals found in the Cap de Creus canyon (A) *Madrepora oculata*, (B) *Lophelia pertusa*, (C) *Desmophyllum dianthus* and (D) *Dendrophyllia cornigera* (source Orejas *et al.*, 2011a).

EXPERIMENTAL WORK: FEEDING ECOLOGY AND GROWTH OF COLD-WATER CORALS

The experimental research developed during the last 5 years focusing on cold-water corals, mainly on feeding ecology and growth rates (Fig. 6), has given new insights into the ecology of these species. These kinds of ecophysiological studies are important to obtain a clearer understanding of suitable locations for their development, their population dynamics, as well as the recovery capacity of these species after disturbances of natural or anthropogenic origin.

Aquaria experiments on capture rates developed for some cold-water corals show how efficient these species are when preying on large zooplankton (Tsounis *et al.*, 2010; Purser *et al.*, 2010). The role of zooplankton as prey on key physiological processes of the cold-water coral species *Desmophyllum dianthus* has also been studied recently, showing that this food source is important in sustaining respiratory metabolism, growth and organic matter release (Nauman *et al.*, 2011). These studies emphasize the importance of zooplankton for these organisms. Furthermore, the role of the cold-water coral communities as a hot spot for carbon recycling has also been proven in some locations (van Oevelen, 2009). Experiments performed in aquaria to study the growth rates of cold-water corals showed that, in some cases, these rates are not as slow as it was previously supposed and documented (Roberts *et al.*, 2009). Recent findings show that, in some cases, their growth rates are even comparable to the growth of some of their tropical counterparts. This is the case for *M. oculata* specimens from the Mediterranean, which showed growth rates comparable to *Galaxea fascicularis* colonies, originally from the Red Sea (Orejas *et al.*, 2011b). Growth rates for the 4 dominant cold-water coral species found in the Cap de Creus canyon showed different values in terms of weight increase, which have been related to possible inter-specific physiological differences (Orejas *et al.*, 2011a). The relative high growth rates recorded for the cold-water coral *M. oculata* suggest a higher recovery potential for these communities after disturbances. Further studies would be important to predict future scenarios, where environmental threats such as ocean acidification could play an important role in cold-water coral growth, as it has already been documented for tropical corals (e.g. Marubini and Atkinson, 1999; Marubini *et al.*, 2008; Schneider and Erez, 2006) and for some cold-water corals (Maier *et al.*, 2009, 2011). Studies on the biology of cold-water coral species are therefore critical to enlarge our knowledge on the ecology of these organisms, providing the basis to develop effective conservation plans and guarantee sustainable management of this community.



Fig. 6:
Growth of *Madrepora oculata* between December 2006 and May 2008, showing the lengthening of branches, new polyps and the covering of the artificial base by new tissue. Colour differences in the last photograph are due to different lighting during photography.

ASSOCIATED ZOOPLANKTON OBSERVATIONS

The video images recorded with the submersible JAGO at about 200 m depth (Fig. 7) show large clouds of Euphausiacea. There are few observations of large swarms of krill near the bottom as observed in the Cap de Creus canyon, which are mainly reported from Antarctic waters (Gutt and Siegel, 1994; Clarke and Tyler, 2008). In the Mediterranean, observations of the most abundant krill species *Meganyctiphanes norvegica* show maximum abundances (0.09 ind/m³) in deep areas, around 120 m depth (Tarling *et al.* 2001). The abundance estimated from zooplankton samples in the Cap de Creus canyon for the most abundant species *Nyctiphanes couchii* shows maximum values of up to 0.24 ind/m³ (Gili *et al.*, 2011). These observations are important because krill is the main food source for many marine organisms, especially cetaceans. Zooplankton is also an important prey in the diet of cold-water corals, as has been determined from biochemical analysis (Duineveld *et al.*, 2004; Kiriakoulakis *et al.*, 2005; Cartier *et al.*, 2009) and direct observations on-site (Mortensen, 2001; Freiwald, 2002).



Fig. 7:

Swarms of krill over a patch of *Madrepora oculata* on the southern flank of the Cap de Creus canyon at 200 m depth.

The presence of fish larvae among the zooplankton communities inside the canyon is also important. The species composition of samples obtained within the canyon includes the presence of larvae of some species of high commercial value, such as hake (*Merluccius merluccius*) (Gili *et al.*, 2011), as well as neritic and mesopelagic species (Fig. 8). The presence of fish larvae suggests how important this particular habitat is as a spawning and nursery area for fish. It is known that cold-water corals act as potential places of refuge, breeding and feeding for many deep-sea species in other regions, including commercially important fish (Husebø *et al.*, 2002; Fossa *et al.*, 2002; D'Onghia *et al.*, 2010). The Gulf of Lion is also a breeding ground for commercially important pelagic species like the anchovy *Engraulis encrasicolus* in spring-summer time and the sardine *Sardina pilchardus* in autumn-winter months (Sabatés *et al.*, 2007).

ANTHROPOGENIC IMPACT

The area of Cap de Creus is characterized as being subjected to considerable maritime traffic, being the connection between the populated and industrialized areas of southern France and Barcelona harbour. It is also regarded as a preferential area for intense fishing activities. Additionally, the anthropogenic impact on the coastal region is exacerbated by discharge from sewage collectors, nautical and recreational activities which intensify during the summer months, and fishing (spearfishing) from recreational boats (Lloret *et al.*, 2008a,b). All these activities, especially fishing, have an important impact on the marine ecosystems and require thorough evaluation.

FISHING IMPACT

Trawl fishery

The Vessel Monitoring System (VMS) provides information regarding vessel code, position, time, speed and direction of the boat to the fishing authorities. In the EU, from 2005 onwards, VMS applies to all large fishing vessels exceeding 15 m in length (EC Regulation No. 2244/2003). Since January 1st 2012, VMS has been extended




Fig. 8:

Fish larvae in zooplankton samples collected inside the Cap de Creus canyon. Commercial species are highlighted.

to all vessels longer than 12 m (EC Regulation No. 1224/2009). Although VMS purposes are primarily for fisheries enforcements, they are currently employed as a unique and independent tool to spatially and temporally allocate the distribution patterns of fishing activity (Gerritsen and Lordan, 2011; Mills *et al.*, 2007; Witt and Godley, 2007; Skaar *et al.*, 2011).

The data processed from the VMS has provided the opportunity to visualize the spatial behavior of the fishing industry and hence study its potential influence on benthic communities. The provisional analysis so far developed shows an apparently higher density of signals in spring, autumn and winter in the southern part of the Cap de Creus canyon, whereas such fishing activity moves towards a northern canyon during the summer (Fig. 9).

Artisanal fishing

Artisanal fisheries have a long tradition in the Cap de Creus area, and are defined by combining fishing arts, target species, fishing geographical zones and seasonality (Table 1). Despite alternating seasons for target species and closed fishing seasons, this activity takes place all year round, weather permitting (Table 1). Fishermen's local knowledge determines this seasonality by considering a species' behavior and its abundance throughout the year (Stelzenmüller *et al.*, 2007). Fishermen tend to state a lack of specificity in the choice of their fishing grounds, but in fact weather and the previous day's fishing experience determine their destination. Despite their activity, a decline in artisanal fisheries has been observed during the last decades to the benefit of progressive expansion of other practices such as semi-industrial fishing and tourism-related activities (Gómez *et al.*, 2006).

In order to define the degree of impact that fishing gears exert on benthic communities, an overlap value was established (Purroy *et al.*, 2010). The assigned values ranged from 0 to 4 depending on the physical coexistence of the main fishing gears (surface and bottom longlines, trammel nets and gillnets). The absence of fishing gears detected in a given area was attributed the value 0, for those areas where only one fishing gear was active the given value

was 1, and so on. A value of 4 was not obtained, indicating that confluence of all 4 gears has not been recorded in the area. The coverage percentage of a particular gear attained values of 60%; the coincidence of two gears corresponded to a value of 29%, whereas interaction of 3 fishing gears being used simultaneously was only observed in 11% of cases. The resulting map reflected the limitations in the coexistence of particular fishing types, i.e. trammel nets and longlines hardly coincide, due to the incompatibility of their fishing techniques.

Comparing Fig. 11, which highlights the well-preserved communities in the study area, and Fig. 10 with the overlap value map, a relationship between the best-preserved communities and the degree of impact of fishing activities can be observed. Thus, areas of high ecological interest coincide with those with less overlap of fishing gears, with values ranging from 0 to 2.

In both cases, artisanal fishing and bottom trawling, the study of the fishing footprint revealed those areas where little fishing gear overlap occurred, coinciding with those areas of high ecological interest, and indicating the degree of impact that fishing gears have on benthic habitats (Gili *et al.*, 2011). The fishing footprint could be used to localize areas where low fishing impact has presumably helped towards the preservation of benthic communities, where potentially pristine areas with high diversity rates can be found. Thus, the fishing footprint appears as an appropriate tool for the assessment and management of areas of interest for conservation.

Fishing impact from ROV images

We investigated the level of the potential anthropogenic impact on cold-water coral species by quantifying the amount of benthic long-line fishing gear and other anthropogenic remnants observed within the canyon (Orejas *et al.*, 2009). Results showed an average of one fishing line every 10 m of video transect. The hard substrate areas snag the long-lines, where they can get entangled by the coral colonies and therefore represent a potential threat to the corals (Fig 12 A, B). Other fishing gears such as gillnets and plastic debris were occasionally observed. Although there is no bottom trawling in the canyon head, the influence of trawling was presumably noticed in the canyon rims where depositions of detritic communities and cold-water coral remains indicated trawling net activity.




Fig. 9:

Distribution map of fishing vessel locations in the Cap de Creus area in different seasons.




Fig. 10:

Overlap value in the study area of Cap de Creus. Minimum value of overlap is 0 when no fishing gear is present, and maximum value is 3 when 3 different gears coexist.




Fig. 11:

Delimitation of communities and the most representative groups in the study area.

MARINE PROTECTED AREA

The geomorphology and oceanography of the Cap de Creus region, as well as the high number of benthic communities hosted in this area, have a decisive influence in converting this region into a remarkable case for the Mediterranean. The relatively small Cap de Creus area holds high species (α) and habitat (γ) diversity, both representatives of most Mediterranean biocenoses. In this area, many coastal communities (i.e. *Posidonia oceanica* beds, *Cymodocea nodosa* beds, and Maërl beds), and shelf-slope communities (i.e. shelf communities dominated by gorgonians, *Leptometra phalangium* beds) are found. At the same time, this region is one of the few known sites in the Mediterranean Sea to

host cold-water coral communities. All of them have been identified as Sensitive Habitats by the European Commission. The shelf and slope region have been proposed as a marine Site of Community Importance (p-SCI) candidate under the Natura 2000 network. The information compiled on the characteristics of the area has been produced through the project LIFE+ INDEMARES "Inventory and designation of marine Natura 2000 areas in the Spanish seas".

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
Trammel nets												
<i>Scorpaena</i> spp.												
Mullidae												
<i>Palinurus elephas</i>												
Soleidae												
<i>Sepia officinalis</i>												
Gillnets												
<i>Sarda sarda</i>												
<i>Diplodus sargus</i>												
Mullidae												
<i>Merluccius merluccius</i>												
<i>Pagellus acarne</i>												
<i>Atherina boyeri</i>												
<i>Pagellus erythrinus</i>												
Gillnets-Trammel nets												
<i>Sarda sarda</i>												
Pots												
<i>Octopus vulgaris</i>												
<i>Sepia officinalis</i>												
Longlines												
<i>Sparus aurata</i>												
<i>Diplodus sargus</i>												
<i>Conger conger</i>												
<i>Pagellus erythrinus</i>												
<i>Phycis blennoides</i>												
<i>Merluccius merluccius</i>												
<i>Dicentrarchus labrax</i>												
Bottom longlines												
<i>Merluccius merluccius</i>												
<i>Pagellus bogaraveo</i>												
Miscellaneous gears												
<i>Paracentrotus lividus</i>												
<i>Mytilus galloprovincialis</i>												
<i>Corallium rubrum</i>												
<i>Nereis</i> spp.												
Handlines and pole-lines												
<i>Loligo vulgaris</i>												
Boat or vessel seines+A14												
<i>Gymnammodytes cicereus</i>												
<i>Donax trunculus</i>												
<i>Acanthocardia tuberculata</i>												

Tab. 1:

Monthly specificity of *métiers* for each fishing gear (<http://www.faocopemed.org>).
The concept of *métier* corresponds to a combination of gear, target species and fishing geographic zone,
and is applied in order to define the real effort invested in a resource.




Fig. 12:

- A) Colonies of *Madrepora oculata* entangled in a long line.
B) Long lines on the vertical walls of the Cap de Creus canyon.

Acknowledgements

The authors thank Fugro Survey, AOA Geophysics and the University of Barcelona for the multibeam bathymetry of the Cap de Creus canyon, the owners of the ROVs (FIELAX, INSTALSUB, Spain and Gavin Newman, UK), as well as the JAGO team (J. Schauer and K. Hissmann) from IFM-GEOMAR (Germany). We also thank the Spanish General Secretariat for the Sea (Ministerio de Agricultura, Alimentación y Medio Ambiente) for the vessel monitoring systems (VMS) data and the Fundación Biodiversidad for their continuous support. Thanks are also due to the crew of the Research Vessel "García del Cid" as well as to the technical personnel of the Marine Technology Unit (UTM, CSIC). The research developed in the Cap de Creus canyon and shelf has been supported by the Spanish National Project DEEPCORAL (CTM2005-07756-C02-02/MAR), the Acciones Complementarias project (CTM2005-24174-E, CTM2006-27063-E/MAR, CTM2007-28758-E/MAR), the European projects HERMES (Goce-CT-2005-511234-I) and HERMIONE (Grant Agreement Number 226354) and the European Life + project INDEMARES.

REFERENCES

- Canals, M., Danovaro, R., Heussner, S., Lykousis, V., Puig, P., Trincardi, P., Calafat, A.M., Durrieu de Madron, X., Palanques, A. and A. Sánchez-Vidal, 2009. Cascades in Mediterranean Submarine Grand Canyons. *Oceanography*, **22**, 26–43.
- Canals, M., Puig, P., de Madron, X.D., Heussner, S., Palanques, A. and J. Fabres, 2006. Flushing submarine canyons. *Nature*, **444**, 354–357.
- Carlier, A., Le Guilloux, E., Olu, K., Sarrazin, J., Mastrototaro, F., Taviani, M., and J. Clavier, 2009. Trophic relationships in a deep Mediterranean cold-water coral bank (Santa Maria di 836 Leuca, Ionian Sea). *Marine Ecology Progress Series*, **397**, 125–137.
- Clarke, A. And P.A. Tyler, 2008. Adult Antarctic krill feeding at abyssal depths. *Current Biology*, **18**(4), 282–285.
- DeGeest, A.L., Mullenbach, B.L., Puig, P., Nittrouer, C.A., Drexler T.M., Durrieu de Madron X. and D.L. Orange, 2008. Sediment accumulation in the western Gulf of Lion, France: the role of Cap de Creus canyon in linking shelf and slope sediment dispersal systems. *Continental Shelf Research*, **28**, 2031–2047.
- Desbruyères, D., Guille, A. and J. Ramos, 1972–1973. Bionomie benthique du plateau continental de la côte catalane espagnole. *Vie et Milieu*, **23**, 335–363.
- D'Onghia, G., Maiorano, P., Sion, L., Giove, A. *et al.*, 2010. Effects of deep-water coral banks on the abundance and size structure of the megafauna in the Mediterranean Sea. *Deep-Sea Research Part II*, **57**, 397–411.
- Duineveld, G.C.A., Lavaleye, M.S.S. and E.M. Berghuis, 2004. Particle flux and food supply to a seamount cold-water coral community (Galicia Bank, NW Spain). *Marine Ecology Progress Series*, **277**, 13–23.
- Durrieu de Madron, X., Zervakis, V., Theocharis, A. and D. Georgopoulos, 2005. Comments on 'Cascades of dense water around the world ocean'. *Progress in Oceanography*, **64**, 83–90.
- EC (European Commission), 2003. Commission Regulation (EC) No. 2244/2003 of 18 December 2003 laying down detailed provisions regarding satellite-based Vessel Monitoring Systems. *Official Journal of the European Union*, **L333**, 17–27.
- EC (European Commission), 2009. Council Regulation (EC) No. 1224/2009 of 20 November 2009 establishing a Community control system for ensuring compliance with the rules of the Common Fisheries Policy, amending other Regulations (EC). *Official Journal of the European Union*, **L343**, 1–50.
- FAO-COPEMED Project. <http://www.faocopemed.org>
- Fosså, J.H., Mortensen, P.B. and D.M. Furevik, 2002. The deepwater coral *Lophelia pertusa* in Norwegian waters: distribution and fishery impacts. *Hydrobiologia*, **471**, 1–12.
- Freiwald, A. 2002. Reef-forming cold-water corals. In: *Ocean margin systems*. Wefer, G., Billett, D., Hebbeln, D., Jørgensen, B.B., Schlüter, M. and T.C.E. van Weering (eds.). Springer, Heidelberg, pp. 365–385.

- Gerritsen, H. and C. Lordan, 2011. Integrating vessel monitoring systems (VMS) data with daily catch data from logbooks to explore the spatial distribution of catch and effort at high resolution. *ICES Journal of Marine Science*, **68**(1), 245-252.
- Gili, J.M., Madurell, T., Requena, S., Orejas, C., Gori, A., Purroy, A., Domínguez, C., Lo Iacono, C., Isla, E., Lozoya, J.P., Carboneras, C., Grinyó, J. and R. Sardá, 2011. *Caracterización física y ecológica del área marina del Cap de Creus. Informe final área LIFE+INDEMARES (LIFE07/NAT/E/000732)*. Instituto de Ciencias del Mar/CSIC (Barcelona). Coordinación: Fundación Biodiversidad, Madrid, 272pp.
- Gómez, S., Lloret, J., Demestre, M., and V. Riera (2006). Declining artisanal fisheries in the Mediterranean: the case of Cap de Creus (NW Mediterranean). *Coastal Management*, **34**(2), 217-232.
- Gutt, J., and V. Siegel, 1994. Benthopelagic aggregations of krill (*Euphausia superba*) on the deeper shelf of the Weddell Sea (Antarctic). *Deep Sea Research Part I*, **41**(1): 169-178.
- Husebø, A., Nottestad, L., Fosså, J.H., Furevik, D.M., and S.B. Jørgensen, 2002. Distribution and abundance of fish in deep-sea coral habitats. *Hydrobiologia*, **471**, 91-99.
- Kiriakoulakis, K., Fisher, E., Wolff, G.A., Freiwald, A., Grehan, A., and J.M. Roberts, 2005. Lipids and nitrogen isotopes of two deep-water corals from the North-East Atlantic: initial results and implications for their nutrition. In: *Cold-water corals and ecosystems*. Friewald, A. and J.M. Roberts (eds.). Springer, Berlin, pp. 715-729.
- Lastras, G., Canals, M., Urgeles, R., Amblas, D., Ivanov, M., Droz, L., Dennielou, B., Fabrès, J., Schoolmeester, T., Akhmetzhanov, A., Orange, D. and A. García-García, 2007. A walk down the Cap de Creus canyon, Northwestern Mediterranean Sea: recent processes inferred from morphology and sediment bedforms. *Marine Geology*, **246**, 176-192.
- Lloret, J., Zaragoza, N., Caballero, D. and V. Riera, 2008a. Biological and socioeconomic implications of recreational boat fishing for the management of fishery resources in the marine reserve of Cap de Creus (NW Mediterranean). *Fisheries Research*, **91**, 252-259.
- Lloret, J., Zaragoza, N., Caballero, D., Font, T., Casadevall, M. and V. Riera, 2008b. Spearfishing pressure on fish communities in rocky coastal habitats in a Mediterranean marine protected area. *Fisheries Research*, **94**, 84-91.
- Lo Iacono, C., Orejas, C., Gori, A., Gili, J.M., Requena, S., Puig, P. and M. Ribó, 2012. The habitats of the Cap de Creus continental shelf and Cap de Creus Canyon, northwestern Mediterranean. In: *Seafloor Geomorphology as Benthic Habitat: GeoHab Atlas of seafloor geomorphic features and benthic habitats*. Harris, P. and E. Baker (eds.). Elsevier, London, pp. 457-469.
- Maier, C., Hegeman, J., Weinbauer, M.G., Gattuso, J.P., 2009. Calcification of the coldwater coral *Lophelia pertusa* under ambient and reduced pH. *Biogeosciences*, **6**, 1671-1680.
- Maier, C., Watremez, M., Taviani, M., Weinbauer, M.G. and J.P. Gattuso, 2011. Calcification rates and the effect of ocean acidification on Mediterranean coldwater corals. *Proceeding of the Royal Society B* doi: 10.1098/rspb.2011.1763.
- Marubini, F., Atkinson, M.J. 1999. Effects of lowered pH and elevated nitrate on coral calcification. *Marine Ecology Progress Series*, **188**, 117-122.
- Marubini, F., Ferrier-Pages, C., Furla, P. and D. Allemand, 2008. Coral calcification responds to seawater acidification: a working hypothesis towards a physiological mechanism. *Coral Reefs*, **27**, 491-499.
- Mills, C., Townsend, S., Jennings, S., Eastwood, P. and C. Houghton, 2007. Estimating high resolution trawl fishing effort from satellite-based vessel monitoring system data. *ICES Journal of Marine Science*, **64**, 248-255.
- Mortensen, P.B., Hovland, M.T., Fossa, J.H. and D.M. Furevik, 2001. Distribution, abundance and size of *Lophelia pertusa* coral reefs in mid-Norway in relation to seabed characteristics. *Journal of the Marine Biological Association of the UK*, **81**, 581-597.
- Naumann, M.S., Orejas, C., Wild, C., and C. Ferrier-Pagès, 2011. First evidence for zooplankton feeding sustaining key physiological processes in a scleractinian cold-water coral. *Journal of Experimental Biology*, **214**, 3570-3576.
- Ogston, A.S., Drexler, T.M. and P. Puig, 2008. Sediment delivery, resuspension, and transport in two contrasting canyon environments in the southwest Gulf of Lion. *Continental Shelf Research*, **28**, 2000-2016.
- Orejas, C., Gori, A., Lo Iacono, C., Puig, P., Gili, J.M. and M.R.T. Dale, 2009. Cold-water corals in the Cap de Creus canyon, northwestern Mediterranean: spatial distribution, density and anthropogenic impact. *Marine Ecology Progress Series*, **397**, 37-51.
- Orejas, C., Ferrier-Pagès, C., Reynaud, S., Gori, A., Beraud, E., Tsounis, G., Allemand, D. and J.M. Gili, 2011a. Long-term growth rates of four Mediterranean cold-water coral species maintained in aquaria. *Marine Ecology Progress Series*, **429**, 57-65.
- Orejas, C., Ferrier-Pagès, C., Reynaud, S., Tsounis, G., Allemand, D., and J.M. Gili, 2011b. Experimental comparison of skeletal growth rates in the cold-water coral *Madrepora oculata* Linnaeus, 1758 and three tropical scleractinian corals. *Journal of Experimental Marine Biology and Ecology*, **405**, 1-5.
- Palanques, A., Durrieu de Madron, X., Puig, P., Fabres, J., Guillén, J., Calafat, A., Canals, M., Heussner, S. and J. Bonnin, 2006. Suspended sediment fluxes and transport processes in the Gulf of Lions submarine canyons. The role of storms and dense water cascading. *Marine Geology*, **234**, 43-61.
- Pérès, J.M. and J. Picard, 1964. Nouveau Manuel de bionomie benthonique de la Mer Méditerranée. *Recueil des Travaux de la Station Marine d'Endoume*, **31** (47), 1-137.

- Pérès, J.M., 1985. History of the Mediterranean Biota and the Colonization of the Depths. In: *Key Environments: Western Mediterranean*. R. Margalef (ed.), Pergamon Press, New York, pp. 198-232.
- Puig, P., Palanques, A., Orange, D.L., Lastras, G. and M. Canals, 2008. Dense shelf water cascades and sedimentary furrow formation in the Cap de Creus Canyon, north-western Mediterranean Sea. *Continental Shelf Research*, **28**, 2017–2030.
- Purser, A., Larsson, A.I., Thomsen, L. and D. van Oevelen, 2010. The influence of flow velocity and food concentration on *Lophelia pertusa* (Scleractinia) zooplankton capture rates. *Journal of Experimental Marine Biology and Ecology*, **395**(1–2), 55-62.
- Purroy, A., Requena, S., Sardá, R. and J.M. Gili, 2010. Spatial assessment and impact of artisanal fisheries' activity in Cap de Creus. In: *Geographic technologies applied to Marine Spatial Planning and Integrated Coastal Zone Management*. Calado, H. and A. Gil (eds.). Centro de Informação Geográfica e Planeamento Territorial, Universidade dos Açores. Ponta Delgada, Portugal, pp. 15-22.
- Roberts, J.M., Wheeler, A.J., Freiwald, A., and S.D. Cairns, 2009. *Cold-Water Corals. The biology and geology of Deep-Sea Coral habitats*. Cambridge University Press, Cambridge, 334pp.
- Sabatés, A., Olivar, M.P., Salat, J., Palomera, I., and F. Alemany, 2007. Physical and biological processes controlling the distribution of fish larvae in the NW Mediterranean. *Progress in Oceanography*, **74**, 355–376.
- Schneider K and J. Erez, 2006. The effect of carbonate chemistry on calcification and photosynthesis in the hermatypic coral *Acropora eurystoma*. *Limnology and Oceanography*, **51**, 1284-1293.
- Skaar, K.L., Jorgensen, T., Ulvestad, B.K.H. and A. Engas, 2011. Accuracy of VMS data from Norwegian demersal stern trawlers for estimating trawled areas in the Barents Sea. *ICES Journal of Marine Science*, **68**, 1615-1620.
- Stelzenmüller, V., Maynou, F. and P. Martin, 2007. Spatial assessment of benefits of a coastal Mediterranean Marine Protected Area. *Biological Conservation*, **136**, 571-583.
- Tarling, G.A., Matthews, J.B.L., David, P., Guerin, O. and F. Buchholz, 2001. The swarm dynamics of northern krill (*Meganyctiphanes norvegica*) and pteropods (*Cavolina inflexa*) during vertical migration in the Ligurian Sea observed by acoustic Doppler current profiler. *Deep-Sea Research I*, **48**, 1671-1686.
- Tsounis, G., Orejas, C., Reynaud, S., Gili, J.M., Allemand, D. and C. Ferrier-Pagès, 2010. Prey capture rates in four Mediterranean cold water corals. *Marine Ecology Progress Series*, **398**, 149–155.
- Ulses, C., Estournel, C., Bonnin, J., Durrieu de Madron, X. and P. Marsaleix, 2008. Impact of storms and dense water cascading on shelf-slope exchanges in the Gulf of Lion (NW Mediterranean). *Journal of Geophysical Research*, **113**, 1-18.
- Van Oevelen, D., Duineveld, G., Lavaleye, M., Mienis, F., Soetaert, K. and C.H.R. Heip, 2009. The cold-water coral community as a hot spot for carbon cycling on continental margins: a food-web analysis from Rockall Bank (northeast Atlantic). *Limnology and Oceanography*, **54**(6), 1829-1844.
- Witt, M. and B. Godley, 2007. A step towards seascape scale conservation: using Vessel Monitoring Systems (VMS) to map fishing activity. *PLoS ONE*, **2** (10), e1111.